Anti-Human CD85j / LILRB1 (GHI/75) Mouse IgG2a Recombinant Antibody, PBS Only




Catalog Number:65623-1-PBS

Basic Information	Catalog Number: 65623-1-PBS	GenBank Accession Number: BC015731	Purification Method: Protein A purification
	Size: 1mg , 2mg/ml	GenelD (NCBI): 10859	CloneNo.: GHI/75
	Source: Mouse Isotype: IgG2a	Full Name: leukocyte immunoglobulin-like	d
		receptor, subfamily B (with TM an ITIM domains), member 1	
		Calculated MW: 71 kDa	
Applications	Tested Applications: FC		
	Species Specificity: human		
Background Information	LILRB1, also named as CD85j or ILT2, includes four C2 Ig domains binding HLA class I molecules in the extracellular region and a cytoplasmic domain containing four immunoreceptor tyrosine-based inhibition motifs (ITIM). LILRB1 is a surface glycoprotein detected on the surface of B cells, monocytes, dendritic cells, T cells and NK cells.		
Storage	Storage: Store at -80°C. Storage Buffer:		


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^{^6} human PBMCs were surface stained with 0.25 ug Anti-Human CD85j / LILRB1 (GHI/75) Mouse IgG2a Recombinant Antibody (65623-1-MR, Clone: GHI/75) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Jg, Clone: C1.18.4) and Coralite®488-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 65623-1-PBS in a different storage buffer



1x10^{^6} human PBMCs were surface stained with 0.25 ug Anti-Human CD85j / LILRB1 (GHI/75) Mouse IgG2a Recombinant Antibody (65623-1-MR, Clone: GHI/75) (red) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clonic C1.18.4) (blue) and CoraLite®488-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 65623-1-PBS in a