For Research Use Only

## Anti-Mouse PD-L1/CD274 (10F.9G2) Mouse IgG2a Recombinant Antibody



Catalog Number: 65685-1-MR

**Basic Information** 

Catalog Number:

65685-1-MR

Size:

100ug , 1000  $\mu g/ml$ Source:

Mouse

Isotype: lgG2a

GenBank Accession Number:

BC066841 GeneID (NCBI): 60533

Full Name:

CD274 antigen

**Purification Method:** 

Protein A purification

CloneNo.: 10F.9G2

Recommended Dilutions:

FC: 0.25 ug per 10^6 cells in a 100  $\mu$ l

suspension

**Applications** 

**Tested Applications:** 

Species Specificity:

mouse

**Positive Controls:** 

FC: Anti-CD3/CD28 treated mouse splenocytes,

## **Background Information**

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner.

Storage


Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH7.3

in USA), or 1(312) 455-8498 (outside USA)

## **Selected Validation Data**



1x10^6 untreated or anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse PD-L1/CD274 (10F.9G2) Mouse IgG2a RecAb (65685-1-MR, Clone:10F.9G2) and PE-conjugated Goat Anti-Mouse IgG. Cells were co-stained with APC Anti-Mouse CD3. Cells were not fixed.



1x10^6 anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse PD-L1/CD274 (10F.9G2) Mouse IgG2a RecAb (65685-1-MR, Clone: 10F.9G2) (red) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue) and PE-conjugated Goat Anti-Mouse IgG. 1x10^6 untreated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse PD-L1/CD274 (10F.9G2) Mouse IgG2a RecAb (65685-1-MR,

