

For Research Use Only

Anti-Mouse PD-L1/CD274 (10F.9G2) Rat IgG2a Recombinant Antibody

Catalog Number: 65685-2-TR



Basic Information

Catalog Number:

65685-2-TR

Size:

100ug, 1000 µg/ml

Source:

Rat

Isotype:

IgG2a

GenBank Accession Number:

BC066841

GeneID (NCBI):

60533

UNIPROT ID:

Q9EP73

Full Name:

CD274 antigen

Purification Method:

Protein A purification

CloneNo.:

10F.9G2

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : Anti-CD3/CD28 treated mouse splenocytes,

Background Information

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner.

Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse PD-L1/CD274 (10F.9G2) Rat IgG2a RecAb (65685-1-TR, Clone: 10F.9G2) and PE anti-rat IgG2a Antibody. Cells were co-stained with APC Anti-Mouse CD3. Cells were not fixed.

1x10⁶ anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse PD-L1/CD274 (10F.9G2) Rat IgG2a RecAb (65685-1-TR, Clone: 10F.9G2) (red) or Rat IgG2a Isotype Control (2A3) (65209-1-Ig, Clone: 2A3) (blue), and PE anti-rat IgG2a Antibody. 1x10⁶ untreated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse PD-L1/CD274 (10F.9G2) Rat IgG2a RecAb (65685-1-TR, Clone: 10F.9G2) (black, dashed) and PE anti-rat IgG2a

