For Research Use Only

NDUFS5 Monoclonal antibody

Catalog Number: 66053-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

66053-1-lg BC001884 Protein A purification
Size: GeneID (NCBI): CloneNo.:

150ul , Concentration: 2400 ug/ml by 4725 CC2A5

Nanodrop and 1053 ug/ml by Bradford UNIPROT ID:

method using BSA as the standard;
O43920

Source:

Recommended Dilutions:
WB 1:1000-1:4000
IHC 1:20-1:200

Mouse NADH dehydrogenase (ubiquinone)
Isotype: Fe-S protein 5, 15kDa (NADHIgG2b coenzyme Q reductase)

Immunogen Catalog Number: Calculated MW:

AG7663 13 kDa

Observed MW: 15 kDa

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HepG2 cells, LO2 cells, HEK-293 cells, fetal human

Purification Method:

brain tissue

IHC: human kidney tissue, human liver tissue

Background Information

NDUFS5 gene is a member of the NADH dehydrogenase (ubiquinone) iron-sulfur protein family. The encoded protein is a subunit of the NADH:ubiquinone oxidoreductase (complex I), the first enzyme complex in the electron transport chain located in the inner mitochondrial membrane. The NDUFS5 mRNA is expressed ubiquitously in human tissues, with a relative higher expression in human heart, skeletal muscle, liver, kidney and fetal heart (PMID: 10070614). The NDUFS5 (15 kDa) subunit located at mitochondrion and contains two C-X9-C motifs that are predicted to form a helix-coil-helix structure, permitting the formation of intramolecular disulfide bonds (PMID: 21310150). Proteinteach's NDUFS5 antibody 66053-1-Ig is capable of detecting the 15 kDa protein in Western blotting.

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 66053-1-1g (NDUFS5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney using 66053-1-Ig(NDUFS5 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney using 66053-1-Ig(NDUFS5 antibody) at dilution of 1:50 (under 40x lens).