For Research Use Only

PMS2 Monoclonal antibody

Catalog Number:66075-1-lg 13 Publications



Purification Method:

CloneNo.:

1G4E6

Protein A purification

Recommended Dilutions:

WB 1:500-1:2000

Basic Information

Catalog Number: GenBank Accession Number:

66075-1-lg BC093921
Size: GeneID (NCBI):

150ul , Concentration: 2700 ug/ml by 5395 Nanodrop and 1580 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; p54278

Source: Full Name:

Mouse PMS2 postmeiotic segregation Isotype: increased 2 (S. cerevisiae)
IgG2a Calculated MW:

IgG2a Calculated MW:

Immunogen Catalog Number: 862 aa, 96 kDa

AG12661 Observed MW:

100 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: human Positive Controls:

WB: A431 cells, HeLa cells

Background Information

PMS2, also named as PMSL2, belongs to the DNA mismatch repair mutL/hexB family. It is a component of the post-replicative DNA mismatch repair system (MMR). It heterodimerizes with MLH1 to form MutL alpha. MulL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. It also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. (PMID: 16873062, PMID: 18206974) Defects in PMS2 are the cause of hereditary non-polyposis colorectal cancer type 4 (HNPCC4). Defects in PMS2 are a cause of mismatch repair cancer syndrome (MMRCS).

Notable Publications

Author	Pubmed ID	Journal	Application
Sachita Ganesa	35388070	Sci Rep	WB
Xin Zhang	38924997	Drug Resist Updat	WB
Zhenwei Yang	38860604	Mol Carcinog	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data





Vairous cells were subjected to SDS PAGE followed by western blot with 66075-1-1g (PMS2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.

A431 cells were subjected to SDS PAGE followed by western blot with 66075-1-lg (PMS2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.