

For Research Use Only

Phospho-TDP43 (Ser403/404) Monoclonal antibody, PBS Only



Catalog Number: 66079-1-PBS

Basic Information

Catalog Number: 66079-1-PBS	GenBank Accession Number: NM_007375	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 23435	CloneNo.: 6B11B12
Source: Mouse	UNIPROT ID: Q13148	
Isotype: IgG2a	Full Name: TAR DNA binding protein	
	Calculated MW: 43 kDa	
	Observed MW: 25 kDa	

Applications

Tested Applications:
WB, FC, Indirect ELISA

Species Specificity:
human, mouse

Background Information

Transactivation response (TAR) DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43) was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major component of the tau-negative and ubiquitin-positive inclusions that characterize amyotrophic lateral sclerosis (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phospho-protein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (PMID: 17023659,19823856, 21666678, 22193176).66079-1-Ig is a mouse monoclonal antibody recognizing TDP-43 only when phosphorylated at 403/404. Immunohistochemical analyses using this antibody only stain the insoluble inclusions in pathologic tissues without normal diffuse nuclear staining.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 66079-1-Ig (Phospho-TDP43 (Ser403/404) antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66079-1-PBS in a different storage buffer formulation.



1×10^6 Jurkat cells were stained with 0.20ug Phospho-TDP43 (Ser403/404) antibody (66079-1-Ig, red) and control antibody (blue). Fixed with 90% MeOH. This data was developed using the same antibody clone with 66079-1-PBS in a different storage buffer formulation.



K-562 cells were subjected to SDS PAGE followed by western blot with 66079-1-Ig (Phospho-TDP43 (Ser403/404) Antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66079-1-PBS in a different storage buffer formulation.