For Research Use Only

Villin Monoclonal antibody

Catalog Number: 66096-1-lg 9 Publications



Basic Information

Catalog Number: GenBank Accession Number:

66096-1-lg BC017303 GeneID (NCBI): Size: 150ul, Concentration: 1000 ug/ml by 7429

Nanodrop and 480 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; P09327

Source: Full Name: Mouse villin 1

Isotype: Calculated MW: lgG1 827aa,93 kDa; 826aa,93 kDa

Immunogen Catalog Number: Observed MW:

AG9637 93-95 kDa **Purification Method:**

Protein A purification

CloneNo.: 2B7B9

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:2500-1:10000 IF/ICC 1:10-1:100

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Human kidney, tissue IP: mouse kidney tissue, IHC: human colon tissue, IF/ICC: HepG2 cells,

Background Information

Villin 1 (VIL1) is a 95-kDa F-actin bundling and severing protein and its expression is restricted to epithelial cells with a brush border, like epithelial cells of the intestinal mucosa, gall bladder, renal proximal tubules and ductuli efferentes of the testis. VIL1 has been reported to be an epithelial cell-specific anti-apoptotic protein, and to have an $important\ function\ in\ regulating\ actin\ dynamics,\ cell\ morphology,\ epithelial-to-mesenchymal\ transitions,\ epithelial-to-mesenchymal\ epithelial-to-mesenchymal\ epithelial-to-mesenchymal\ epithelial-to-mesenchymal\ epithelial-to-mesenchymal\ epithelial-to-mesenchymal\ epithe$ migration and cell survival. In addition, VIL1 is a useful diagnostic marker for of various cancer, like cervical and endometrial adenocarcinomas, renal cell carcinoma. VIL1 was recently identified as a novel biomarker predictive for postoperative recurrence and poorer prognosis of high serum AFP associated HCC.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhang-Mei Peng	25337239	Int J Clin Exp Pathol	IHC
Shanshan Huang	35620578	Oxid Med Cell Longev	IHC
Serika Motoike	33712280	J Pharmacol Sci	IF, IHC

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human kidney tissue were subjected to SDS PAGE followed by western blot with 66096-1-1g (Villin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immun of luorescent analysis of HepG2 cells using 66096-1-lg (Villin antibody) at dilution of 1:25 and Rhodamine-Goat anti-Mouse lgG.



IP result of anti-Villin (IP:66096-1-Ig, 4ug; Detection:66096-1-Ig 1:1000) with mouse kidney tissue lysate 6000ug.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66096-1-Ig (Villin antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66096-1-Ig (Villin antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).