For Research Use Only

IL-6 Monoclonal antibody, PBS Only

Catalog Number:66146-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

66146-1-PBS

GeneID (NCBI):

BC015511

100ug, Concentration: 1mg/ml by

CloneNo.: 1A3B4

Nanodrop:

ENSEMBL Gene ID: ENSG00000136244

Mouse Isotype:

lgG2b

UNIPROT ID: P05231

Full Name:

24 kDa

Immunogen Catalog Number:

interleukin 6 (interferon, beta 2)

AG10646

Calculated MW: 212 aa, 24 kDa Observed MW:

Applications

Tested Applications: WB, IF/ICC, ELISA

Species Specificity:

human

Background Information


 $Interleukin-6 (IL-6) is an interleukin that acts as both a pro-inflammatory and anti-inflammatory cytokine. \ IL-6 in the control of the co$ protein is secreted by a variety of cell types including T cells and macrophages as phosphorylated and variably glycosylated molecule. IL-6 plays an essential role in the final differentiation of B-cells into Ig-secreting cells involved in lymphocyte and monocyte differentiation. It induces myeloma and plasmacytoma growth and induces nerve cells differentiation Acts on B-cells, T-cells, hepatocytes, hematopoietic progenitor cells and cells of the CNS. IL-6 is also considered a myokine, a cytokine produced from muscle, and is elevated in response to muscle contraction. IL-6 has been shown to interact with interleukin-6 receptor and glycoprotein 130. Additionally, IL-6 is involved in hematopoiesis, bone metabolism, and cancer progression, and has been defined an essential role in directing transition from innate to acquired immunity.

Storage


Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data



Non-treated and protein transport inhibitor treated U-87 MG cells were subjected to SDS PAGE followed by western blot with 66146-1-lg (IL-6 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66146-1-PBS in a different storage buffer formulation.



Non-treated, LPS and protein transport inhibitor treated HUVEC cells were subjected to SDS PAGE followed by western blot with 66146-1-lg (IL-6 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66146-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed LPS and Brefeldin A treated HUVEC cells using IL-6 antibody (66146-1-Ig, Clone: 1A3B4) at dilution of 1:800 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 66146-1-PBS in a different storage buffer formulation.