For Research Use Only

NSE Monoclonal antibody, PBS Only

Catalog Number:66150-1-PBS

Featured Product



Basic Information

Catalog Number: 66150-1-PBS Size: 100ug , Concentration: 1mg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG19106 GenBank Accession Number: BC002745 GeneID (NCBI): 2026 UNIPROT ID: P09104 Full Name: enolase 2 (gamma, neuronal) Calculated MW: 47 kDa Observed MW: 47 kDa

Purification Method: Protein A purification CloneNo.: 6F8G3

Applications

Tested Applications: WB, IHC, IF-P, FC (Intra), Indirect ELISA Species Specificity: human, mouse, rat, pig

Background Information

NSE, also named as ENO2, belongs to the enolase family. Enolases are cytoplasmic glycolytic enzymes that may be involved in differentiation. The enolase has three isoenzymes, alpha, beta and gamma. The alpha form is expressed in most tissues, whereas the beta form is expressed in muscle tissue. The gamma enolase (ENO2), a homodimer, is primarily localized in neurons and neuroendocrine cells and is a cancer diagnostic marker for brain tumors (PMID:7520111). ENO2 plays a role in the glycolysis-related energy pathway and might be involved in higher metabolic activity during the day than at night, at least in part.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66150-1-lg (NSE antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.




1X10⁶ SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human NSE (66150-1-1g, Clone:6F8G3) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.




WB result of NSE antibody (66150-1-Ig; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NSE transfected U-251 cells. This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66150-1-Ig (NSE antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66150-1-Ig (NSE antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using NSE antibody (66150-1-lg, Clone: 6F8G3) at dilution of 1:400 and Coralite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using NSE antibody (66150-1-lg, Clone: 6F8G3) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66150-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 66150-1-lg (NSE antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed

using the same antibody clone with 66150-1-PBS in

a different storage buffer formulation.