

STMN3 Monoclonal antibody

Catalog Number: 66152-1-Ig

Basic Information

Catalog Number: 66152-1-Ig	GenBank Accession Number: BC009381	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop and 547 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 50861	CloneNo.: 1F12B1
Source: Mouse	UNIPROT ID: Q9NZ72	Recommended Dilutions: WB 1:500-1:2000
Isotype: IgG2b	Full Name: stathmin-like 3	
Immunogen Catalog Number: AG18844	Calculated MW: 21 kDa	
	Observed MW: 21 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human brain tissue,
Species Specificity: human	

Background Information

Stathmin family phosphoproteins participate in the control of microtubule dynamics and have been proposed to be involved in the control of neuronal differentiation. Neuron specific Stathmin-3 (STMN3) is a novel STAT3 (signal transducer and activator of transcription 3) interacting protein. STAT3 is a key contributor to cancer cell migration and invasion. STMN3 exhibits microtubule-destabilizing activity, which is antagonized by STAT3. SCG10 is a widely studied and recognized neuronal differentiation marker, STMN3 is a SCG10-like protein (SCLIP) and is present from the earliest stages of hippocampal neuron differentiation in culture at vesicle-like structures following dynamic microtubules. RNAi mediated inhibition of this gene resulted in increased axonal branching.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 66152-1-Ig (STMN3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.