For Research Use Only

Vitamin D binding protein Monoclonal antibody, PBS Only



Catalog Number: 66175-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

66175-1-PBS

GeneID (NCBI):

Protein G purification

BC057228

CloneNo.:

100ug, Concentration: 1mg/ml by Nanodrop:

1E4D10

Mouse

UNIPROT ID: P02774

Isotype: lgG1

Full Name: group-specific component (vitamin D

binding protein)

Immunogen Catalog Number: AG9803

Calculated MW:

474 aa, 53 kDa Observed MW:

52-58 kDa

Applications

Tested Applications:

Indirect ELISA, IHC, WB

Species Specificity:

Background Information

Vitamin D binding protein is a sparsely glycosylated serum protein responsible for highly specific binding and tissue-specific delivery of vitamin D and its metabolites. In addition, it is also an actin scavenger, and is the precursor to the immunomodulatory protein, Gc-MAF. Vitamin D binding protein has been proposed to have significant roles in C5a chemotaxis, osteoclast development and possibly in macrophage activation/recruitment.


Storage

Storage:


Store at -80°C. Storage Buffer:

PBS Only

Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 66175-1-1g (Vitamin D binding protein antibody at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66175-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66175-1-lg (Vitamin D binding protein antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66175-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66175-1-1g (Vitamin D binding protein antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66175-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66175-1-Ig (Vitamin D binding protein antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66175-1-PBS in a different storage buffer formulation.



1X10^6 U-937 cells were intracellularly stained with 0.4 ug Anti-Human Vitamin D binding protein (66175-1-lg, Clone:1E4D10) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse I gG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66175-1-PBS in a different storage buffer formulation.