

## CD71 Monoclonal antibody

Catalog Number: 66180-1-Ig

Featured Product

28 Publications

## Basic Information

## Catalog Number:

66180-1-Ig

## Size:

150ul, Concentration: 1500 ug/ml by Nanodrop;

## Source:

Mouse

## Isotype:

IgG2a

## Immunogen Catalog Number:

AG21612

## GenBank Accession Number:

BC001188

## GeneID (NCBI):

7037

## ENSEMBL Gene ID:

ENSG00000072274

## UNIPROT ID:

P02786

## Full Name:

transferrin receptor (p90, CD71)

## Calculated MW:

85 kDa

## Observed MW:

90 kDa

## Purification Method:

Protein A purification

## CloneNo.:

3C11F11

## Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:2000-1:8000

IF-P 1:200-1:800

IF/ICC 1:200-1:800

## Applications

## Tested Applications:

WB, IHC, IF/ICC, IF-P, IP, ELISA

## Cited Applications:

WB, IHC, IF, Blocking assay

## Species Specificity:

human

## Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB: U-251 cells, A549 cells, HeLa cells, Raji cells, TF-1 cells, U2OS cells, A431 cells, HepG2 cells, K-562 cells

IP: HeLa cells,

IHC: human placenta tissue, human breast cancer tissue

IF-P: human breast cancer tissue, MCF-7 cells

IF/ICC: MCF-7 cells, K-562 cells

## Background Information

CD71, also known as transferrin receptor protein 1 (TfR1), is a transmembrane glycoprotein composed of two disulfide-linked monomers, each of 90 kDa molecular weight. Each monomer binds one holo-transferrin molecule creating an iron-Tf-TfR complex which enters the cell by endocytosis. CD71 is present on actively proliferating cells and is essential for iron transport into proliferating cells.

## Notable Publications

Author	Pubmed ID	Journal	Application
Ping Han	36309297	Free Radic Biol Med	WB
Lingjuan Wang	34732689	Cell Death Discov	WB
Yangli Tian	34081703	PLoS Genet	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66180-1-Ig (CD71 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.




WB result of CD71 antibody (66180-1-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD71 transfected HeLa cells.




Untreated and PNGase F-treated lysates of Raji cells and HeLa cells were subjected to SDS PAGE followed by western blot with 66180-1-Ig (CD71 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 66180-1-Ig (CD71 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 66180-1-Ig (CD71 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CD71 antibody (66180-1-Ig, Clone: 3C11F11) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).





Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using CD71 antibody (66180-1-Ig, Clone: 3C11F11) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IP result of anti-CD71 (IP:66180-1-Ig, 5ug; Detection:66180-1-Ig 1:1000) with HeLa cells lysate 2400ug.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 66180-1-Ig (CD71 antibody) at dilution of 1:40000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed K-562 cells using CD71 antibody (66180-1-Ig, Clone: 3C11F11 ) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).

Immunofluorescent analysis of (4% PFA) fixed K-562 cells using CD71 antibody (66180-1-Ig, Clone: 3C11F11 ) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).