For Research Use Only

IL-23 p19 Monoclonal antibody

Catalog Number:66196-1-lg 6 Publications



Basic Information

Catalog Number:

GenBank Accession Number: BC066268

Purification Method:

66196-1-lg

GeneID (NCBI):

Protein G purification

Size: 150ul, Concentration: 1500 ug/ml by 51561

CloneNo.: 1D5D3

Nanodrop and 779 ug/ml by Bradford UNIPROT ID:

Recommended Dilutions:

method using BSA as the standard;

O9NPF7

IHC 1:50-1:500

Source:

Full Name:

interleukin 23, alpha subunit p19

Mouse Isotype: lgG1

Calculated MW:

189 aa, 21 kDa

Immunogen Catalog Number:

AG19406

Applications

Tested Applications:

IHC FIISA

Cited Applications:

WB, IHC, IF, ELISA

Species Specificity: human

Cited Species:

human, mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human stomach cancer tissue, human pancreas

Background Information

Interleukin 23 (IL-23) is a member of the IL12 cytokine family and composed of two subunits, IL12p40 and IL23p19. It is produced by antigen presenting cells and has been shown to promote the production and survival of a distinct lineage of T-cells called TH17 cells. A functional receptor for IL-23 (the IL-23 receptor) has been identified and is $composed \ of \ Il-12R\beta 1 \ and \ Il-23R. \ IL-23 \ is \ expressed \ chiefly \ by \ the \ macrophages \ and \ DCs. \ The \ Il-23R \ is \ found \ on \ and \ DCs.$ memory T cells, NKT cells, macrophages, DCs, and naive T cells upon activation by TGF-βand IL-6. The main biological effects of IL-23 identified initially consist of stimulation of antigen presentation by DCs, T cell differentiation to Th17 cells, and production of interferon-y (IFN-y). IL-23 acts also as an end-stage effector cytokine through direct action on macrophages.

Notable Publications

Author	Pubmed ID	Journal	Application
Teresa W-M Fan	36150727	J Immunol	
YuanPing Hai	36128805	Thyroid	IHC
Rui Sun	31097538	Gut	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 66196-1-lg (IL-23A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 66196-1-lg (IL-23A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).