For Research Use Only

CHD4 Monoclonal antibody

Catalog Number:66222-1-lg

Featured Product

1 Publications



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number: BC038596

66222-1-lg Protein G purification GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 1300 ug/ml by 1108 3D9G2 Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} Recommended Dilutions:

method using BSA as the standard; Q14839 WB 1:500-1:2000 Source: IHC 1:50-1:500 Full Name: chromodomain helicase DNA binding IF/ICC 1:50-1:500 Mouse

Isotype: protein 4

lgG1 Calculated MW: Immunogen Catalog Number: 1937 aa. 221 kDa AG5729 Observed MW:

260 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MCF-7 cells, HEK-293 cells IHC: human cervical cancer tissue,

IF/ICC: HEK-293 cells,

Background Information

Chromodomain helicase DNA-binding protein 4 (CHD4) is an ATP dependent chromatin remodeler and a major subunit of the repressive NuRD complex [PMID:9790534]. This complex plays key roles in transcriptional regulation and reorganization and maintenance of chromatin structure, as it targets pericentromeric heterochromatin and has recently been implicated in DNA damage repair, being rapidly recruited to DNA double strand breaks $[PMID: 17694084]. \ CHD4 is a novel BRIT1 binding partner that regulates the HR repair process, which is a critical and the process of the$ pathway of repairing DNA double strand breaks and plays an essential role in maintaining genomic integrity [PMID:22219182]. The calcualted molecular weight of CHD4 is 218 kDa, but modified CHD4 is about 250-280 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenxiu Dai	38272027	Dev Cell	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




WB result of CHD4 antibody (66222-1-lg, 1:1000) with si-Control and si-CHD4 transfected HEK293 cells.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 66222-1-Ig (CHD4 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



MCF-7 cells were subjected to SDS PAGE followed by western blot with 66222-1-1g (CHD4 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (10% Formaldehyde) fixed HEK-293 cells using 66222-1-Ig(CHD4 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated Goat Anti-Mouse IgG(H+L).