

For Research Use Only

HPSE Monoclonal antibody

Catalog Number: 66226-1-Ig

Featured Product

3 Publications



Basic Information

Catalog Number: 66226-1-Ig	GenBank Accession Number: BC051321	Purification Method: Protein G purification
Size: 150ul, Concentration: 4800 µg/ml by Nanodrop and 2080 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 10855	CloneNo.: 1D8B8
Source: Mouse	Full Name: heparanase	Recommended Dilutions: WB 1:500-1:2000 IHC 1:200-1:1000 IF 1:50-1:500
Isotype: IgG1	Calculated MW: 543 aa, 61 kDa	
Immunogen Catalog Number: AG10067	Observed MW: 50 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells,

IHC : human liver cancer tissue,

IF : HepG2 cells,

Background Information

HPSE (Heparanase) is also named as HEP, HPA, HPA1, HPR1, HPSE1, HSE1 and belongs to the glycosyl hydrolase 79 family. It is an endoglycosidase that cleaves heparan sulfate proteoglycans (HSPGs) into heparan sulfate side chains and core proteoglycans. HPSE is essential in the disassembly of the extracellular matrix (ECM) by invading cells. It has 3 isoforms produced by alternative splicing with the molecular weight of 61 kDa, 55 kDa and 53 kDa. The full length protein has six glycosylation sites. The cleavage of the 65 kDa form leads to the generation of a linker peptide, and 8 kDa and 50 kDa products. The active form, the 8/50 kDa heterodimer, is resistant to degradation and glycosylation of the 50 kDa subunit appears to be essential for its solubility.

Notable Publications

Author	Pubmed ID	Journal	Application
Rémi Cousin	34677445	Mar Drugs	WB
Xue Liu	31001480	Front Oncol	WB, IHC, IF
Li Li	38334603	Cells	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 66226-1-Ig (HPSE Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66226-1-Ig (HPSE Antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66226-1-Ig (HPSE Antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 66226-1-Ig(HPSE antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).