For Research Use Only

UCH-L1/PGP9.5 Monoclonal antibody, PBS Only



Catalog Number: 66230-1-PBS

Featured Product

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

66230-1-PBS

BC000332

Protein A purification

Size:

GeneID (NCBI):

CloneNo.:

100ug, Concentration: 1mg/ml by

7345

1C9E11

Nanodrop;

UNIPROT ID: P09936

Source Mouse

Isotype

ubiquitin carboxyl-terminal esterase

lgG1

L1 (ubiquitin thiolesterase)

Immunogen Catalog Number: AG6547

Calculated MW: 25 kDa

Observed MW:

27 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat, pig

Background Information


Ubiquitin C-terminal hydrolase L1 (UCHL1) was originally identified as a neuronal protein that accounts for nearly 2% of total brain proteins. UCHL1 activity protects neurons from hypoxic injury, and binding of stroke-induced reactive lipid species to the cysteine 152 (C152) of UCHL1 unfolds the protein and disrupts its function. Reduced hydrolytic activity of mutant UCHL1 is implicated in the pathophysiologic process of Parkinson's and Alzheimer's disease due to abnormal neurotoxic protein aggregation. (PMID: 31356902, PMID: 30760601)

Storage


Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66230-1-Ig (UCHL1 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control. This data was developed using the same antibody clone with 66230-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 66230-1-lg (UCHL1/PGP9.5 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66230-1-PBS in a different storage buffer formulation.



1x10^6 Y79 cells were intracellularly stained with 0.4 ug UCHL1/PGP9.5
Monoclonal antibody (66230-1-Ig, Clone:1C9E11) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same





Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using UCH-L1/PGP9.5 antibody (66230-1-Ig, Clone: 1C9E11) at dilution of 1:800 and Coralite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 66230-1-PBS in a different storage buffer formulation.