

For Research Use Only

PD-L1/CD274 Monoclonal antibody

Catalog Number: 66248-1-Ig

Featured Product

397 Publications



Basic Information

Catalog Number: 66248-1-Ig	GenBank Accession Number: BC074984	Purification Method: Protein G purification
Size: 150ul , Concentration: 2000 ug/ml by Nanodrop;	GeneID (NCBI): 29126	CloneNo.: 2B11D11
Source: Mouse	UNIPROT ID: Q9NZQ7	Recommended Dilutions: WB: 1:2000-1:10000 IHC: 1:5000-1:20000 IF-P: 1:400-1:1600 IF/ICC: 1:50-1:500
Isotype: IgG1	Full Name: CD274 molecule	
Immunogen Catalog Number: AG12443	Calculated MW: 290 aa, 33 kDa	
	Observed MW: 45-50 kDa, 33 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA	Positive Controls: WB : A375 cells, human placenta tissue, pig lung tissue, human skeletal muscle tissue, HepG2 cells, THP-1 cells, RAW 264.7 cells, A549 cells, K-562 cells, HSC-T6 cells IHC : human tonsillitis tissue, human heart tissue, human lung cancer tissue, human placenta tissue, mouse heart tissue IF-P : human placenta tissue, IF/ICC : HeLa cells,
Cited Applications: WB, IHC, IF, IP, CoIP, ChIP	
Species Specificity: human, mouse, rat, pig	
Cited Species: human, mouse, rat, pig	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058).

Notable Publications

Author	Pubmed ID	Journal	Application
Jiacheng Huang	34650926	Front Oncol	IHC
Youqiong Ye	32988398	Genome Med	WB
Hao Zhang	36136350	Brief Bioinform	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human placenta tissue were subjected to SDS PAGE followed by western blot with 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Untreated and IFN gamma treated A375 cells and A549 cells were subjected to SDS PAGE followed by western blot with 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:10000 (under 4x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:10000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 66248-1-Ig(PD-L1/CD274 antibody) at dilution of 1:300 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human placenta tissue using PD-L1/CD274 antibody (66248-1-Ig, Clone: 2B11D11) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).