### For Research Use Only

# PHOX2B Monoclonal antibody

Catalog Number:66254-1-lg 2 Publications



**Purification Method:** 

Recommended Dilutions:

WB 1:1000-1:5000 IHC 1:20-1:200

IF 1:200-1:800

**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number:

66254-1-lg BC017199 Protein A purification Size: GeneID (NCBI): CloneNo.: 1F5F5

32-36 kDa

150ul, Concentration: 1500 µg/ml by 8929 Nanodrop and 1000 µg/ml by Bradford<sub>Full Name</sub>:

method using BSA as the standard; paired-like homeobox 2b

Calculated MW: Mouse 31 kDa Isotype: Observed MW: IgG2a

Immunogen Catalog Number:

AG13196

Species Specificity: human, mouse

**Tested Applications: Positive Controls:** IF THE WR FLISA WB: Neuro-2a cells,

**Cited Applications:** IHC: neuroblastoma tissue. IF: SH-SY5Y cells, Neuro-2a cells

**Cited Species:** human, mouse Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

# **Background Information**

Paired mesoderm homeobox protein 2B (PHOX2B), also named as Neuroblastoma Phox, is a 314 amino acid protein, which contains one homeobox DNA-binding domain and belongs to the paired homeobox family. PHOX2B localizes in the nucleus and is expressed in neuroblastoma, brain and adrenal gland. PHOX2B plays a role in the development of several major noradrenergic neuron populations, including the locus coeruleus. PHOX2B has an association with congenital central hypoventilation syndrome.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yunlong Tao	37974010	Nat Biotechnol	IF
Ouranio Anastasiou	32098802	Sci Signal	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Neuro-2a cells were subjected to SDS PAGE followed by western blot with 66254-1-lg (PHOX2B Antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded neuroblastoma tissue slide using 66254-1-1g (PHOX2B antibody) at dilution of 1:300 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using PHOX2B antibody (66254-1-lg, Clone: 1F5F5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using PHOX2B antibody (66254-1-1g, Clone: 1F5F5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).