#### For Research Use Only

# BRCA1 Monoclonal antibody

Catalog Number:66267-1-lg 1 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein G purification

66267-1-lg Size:

BC115037 GeneID (NCBI):

150ul, Concentration: 1000 µg/ml by 672

CloneNo.: 20D7G6

Bradford method using BSA as the

Calculated MW:

standard:

breast cancer 1, early onset

Mouse Isotype:

1863 aa, 208 kDa Observed MW:

207 kDa

lgG1 Immunogen Catalog Number:

AG19178

**Applications** 

**Cited Applications:** 

Species Specificity:

human

**Tested Applications:** 

human

**Cited Species:** 

## **Background Information**

RCA1, also named as RNF53, plays a central role in DNA repair by facilitating cellular response to DNA repair. It is required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. BRCA1 acts by mediating ubiquitin E3 ligase activity that is required for its tumor suppressor function. It is involved in transcriptional regulation of P21 in response to DNA damage. BRCA1 is required for FANCD2 targeting to sites of DNA damage. It may function as a  $transcriptional\ regulator.\ BRCA1\ inhibits\ lipid\ synthesis\ by\ binding\ to\ inactive\ phosphorylated\ ACACA\ and$ preventing its dephosphorylation. The antibody is specific to BRCA1. BRCA1 appears to produce multiple splice variants. BRCA1 is a nuclear protein with a molecular mass of 220 kDa. The present study describes the isolation and expression of two cDNAs of BRCA1, including a splice variant designated BRCA1D672-4095. BRCA1D672-4095 is generated by exclusion of exon 11 by in-frame splicing and produces a 97 kDa protein. In contrast to BRCA1, BRCA1D672-4095 localizes to the cytoplasm.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Yang Yu	31636387	Oncogene	IF

### Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

Selected Validation Data