For Research Use Only

SDCCAG8 Monoclonal antibody, PBS Only



Catalog Number: 66284-1-PBS

Basic Information

Catalog Number:

66284-1-PBS

100ug, Concentration: 1 mg/ml by

Nanodrop;

Mouse

Isotype:

lgG2b

Immunogen Catalog Number: AG4264

Observed MW:

83 kDa

Purification Method:

Protein A purification CloneNo.:

2H1A4

Applications

Tested Applications: WB, IHC, Indirect ELISA

Species Specificity:

Background Information

SDCCAG8, also named as CCCAP and NY-CO-8, plays a role in the establishment of cell polarity and epithelial lumen formation. It may play a role in ciliogenesis. Loss of SDCCAG8 function as a cause of a retinal-renal $ciliopathy\ and\ validates\ exome\ capture\ analysis\ for\ broadly\ heterogeneous\ single-gene\ disorders.\ SDCCAG8\ is$ localized at both centrioles and interacts directly with OFD1 (oral-facial-digital syndrome 1), which is associated with NPHP-RC.

GenBank Accession Number:

serologically defined colon cancer

BC032454

10806

GeneID (NCBI):

UNIPROT ID: Q86SQ7

Full Name:

antigen 8

Calculated MW: 713 aa, 83 kDa


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 66284-1-1g (SDCCAG8 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66284-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66284-1-lg (SDCCAG8 Antibody) at dilution of 1:200 (under 10x lens). This data was developed using the same antibody clone with 66284-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66284-1-lg (SDC CAG8 Antibody) at dilution of 1:200 (under 40x lens). This data was developed using the same antibody clone with 66284-1-PBS in a different storage buffer formulation.