

For Research Use Only

DAO Monoclonal antibody

Catalog Number: 66296-1-Ig



Basic Information

Catalog Number: 66296-1-Ig	GenBank Accession Number: BC029057	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 1610	CloneNo.: 1C12F4
Source: Mouse	Full Name: D-amino-acid oxidase	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:100-1:400
Isotype: IgG1	Calculated MW: 347 aa, 39 kDa	
Immunogen Catalog Number: AG6928	Observed MW: 39 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, pig, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : pig brain tissue, fetal human brain tissue, pig cerebellum tissue, rat brain tissue, rat cerebellum tissue

IHC : human gliomas tissue, mouse cerebellum tissue

Background Information

D-amino acid oxidase (DAO), with MW of 39 kDa, is a flavoprotein which uses flavin adenine dinucleotide as its prosthetic group. Its substrates include a wide variety of D-amino acids, but it is inactive on the naturally occurring L-amino acids. DAO may play a role in the pathophysiology of schizophrenia and act as a detoxifying agent which removes D-amino acids that accumulate during aging. DAO exists in solution as a mixture of monomer-dimer (80 kDa) and even higher molecular aggregates in equilibrium (170 kDa), depending on the enzyme concentration and forms (PMID: 11130179/PMID: 9210639).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66296-1-Ig (DAO antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 66296-1-Ig (DAO Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieved with Sodium Citrate buffer (pH 6.0).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 66296-1-Ig (DAO Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieved with Sodium Citrate buffer (pH 6.0).