For Research Use Only

## CD47 Monoclonal antibody

Catalog Number:66304-1-lg Featured Product 17 Publications





Basic Information	Catalog Number: 66304-1-1g	GenBank Accession Number: BC010016	Purification Method: Protein A purification			
	Size:	GenelD (NCBI):	CloneNo.:			
	150ul , Concentration: 1500 ug/ml by Nanodrop; Source: Mouse Isotype: IgG2b	961	1E1D8			
		UNIPROT ID: Q08722 Full Name: CD47 molecule Calculated MW:	Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:200-1:800			
				Immunogen Catalog Number: AG14132	323 aa, 35 kDa	
					Observed MW: 50-55 kDa	<i>N</i> :
				Applications	Tested Applications:	Positive
	WB, IF/ICC, ELISA	WB: SKOV-3 cells, A549 cells, HT-1080 cells, Jurkat				
	Cited Applications: WB, IF		I-H1299 cells, HUVEC cells, K-562 cells, TF-1 nan placenta tissue, HeLa cells, HaCaT cells,			
Species Specificity:		cells, MDA-MB-231 cells				
human	IF/ICC : A	A549 cells,				
Cited Species: human, mouse, rat						
Background Information	a five-pass transmembrane attachme nonhematopoietic cells. CD47 interac thrombospondin (THBS1). It is involve	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa			
	a five-pass transmembrane attachme nonhematopoietic cells. CD47 interac thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that rest	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa			
	a five-pass transmembrane attachme nonhematopoietic cells. CD47 interact thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that resu Author Pub	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr ults in the inhibition of phagocyto	s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa osis. Application			
	a five-pass transmembrane attachmenonhematopoietic cells. CD47 interact thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that rest Author Pub Hongying Chen 347	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr ults in the inhibition of phagocyto med ID Journal	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesior otein-a (SIRPa). Upon binding CD47, SIRPa isis. Application logy WB			
	a five-pass transmembrane attachmenonhematopoietic cells. CD47 interact thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that resur- Author Pub Hongying Chen 347 Sukhbir Kaur 348	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr ults in the inhibition of phagocyto med ID Journal 02291 J Nanobiotechnol	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa isis. Application logy WB			
Notable Publications	a five-pass transmembrane attachmenonhematopoietic cells. CD47 interact thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that rest Author Pub Hongying Chen 347 Sukhbir Kaur 348 Jing Li 355	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr ults in the inhibition of phagocyto med ID Journal 02291 J Nanobiotechnol 41476 J Cell Commun S	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa osis. Application logy WB ignal IF,FC			
Notable Publications	a five-pass transmembrane attachmenonhematopoietic cells. CD47 interact thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that resu Author Pub Hongying Chen 347 Sukhbir Kaur 348 Jing Li 355 Storage: Storage: Storage Buffer:	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr ults in the inhibition of phagocyto med ID Journal 202291 J Nanobiotechnol 241476 J Cell Commun S 39950 RSC Adv er shipment.	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa osis. Application logy WB ignal IF,FC			
Background Information Notable Publications Storage	a five-pass transmembrane attachmenonhematopoietic cells. CD47 interact thrombospondin (THBS1). It is involve and migration. CD47 also functions a initiates a signaling cascade that rest Author Pub Hongying Chen 347 Sukhbir Kaur 348 Jing Li 355 Storage: Store at -20°C. Stable for one year aft	ent. CD47 is heavily glycosylated cts with several membrane integr ed in a range of cellular processes s a ligand for signal regulatory pr ults in the inhibition of phagocyto med ID Journal 202291 J Nanobiotechnol 41476 J Cell Commun S 39950 RSC Adv er shipment.	and widely expressed by hematopoietic and ins and also acts a receptor for s, including apoptosis, proliferation, adhesion otein-a (SIRPa). Upon binding CD47, SIRPa osis. Application logy WB ignal IF,FC			

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 66304-1-1g (CD47 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using CD47 antibody (66304-1-Ig, Clone: 1E1D8) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).

WB result of CD47 antibody (66304-1-lg; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD47 transfected A549 cells.