For Research Use Only

IL-21R Monoclonal antibody

Catalog Number: 66319-1-Ig



Purification Method:

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:50-1:200

Basic Information

Catalog Number: GenBank Accession Number:

66319-1-Ig BC004348 Protein A purification
Size: GeneID (NCBI): CloneNo.:
150ul , Concentration: 1500 µg/ml by 50615 1D1C2

Bradford method using BSA as the standard:

Source: Calculated MW:
Mouse 59 kDa

Isotype: Observed MW: IgG2b 55-65 kDa

Immunogen Catalog Number:

AG21295

Positive Controls:

WB: HepG2 cells, Jurkat cells, human placenta tissue,

K-562 cells, TF-1 cells, MOLT-4 cells

IHC: human tonsillitis tissue,

Applications

Tested Applications: IHC, WB,ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IL21R, a receptor for interleukin-21, belongs to the type I cytokine receptor family. IL21R has been shown to form a heterodimeric receptor complex with the common gamma-chain, a receptor subunit also shared by the receptors for interleukin 2, 4, 7, 9, and 15. This receptor transduces the promoting signal of IL21, and is important for the proliferation and differentiation of T cells, B cells, and natural killer (NK) cells. The ligand binding of this receptor leads to the activation of multiple downstream signaling molecules, including JAK1, JAK3, STAT1, and STAT3.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66319-1-lg (IL-21R antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66319-1-Ig (IL21R antibody) at dilution of 1:100 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66319-1-Ig (IL21R antibody) at dilution of 1:100 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).