

For Research Use Only

TH Monoclonal antibody

Catalog Number: 66334-1-Ig

Featured Product

25 Publications



Basic Information

Catalog Number:

66334-1-Ig

Size:

150ul, Concentration: 2000 ug/ml by Nanodrop and 806 ug/ml by Bradford method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG23075

GenBank Accession Number:

BC104967

GeneID (NCBI):

7054

UNIPROT ID:

P07101

Full Name:

tyrosine hydroxylase

Calculated MW:

528 aa, 59 kDa

Observed MW:

55 kDa

Purification Method:

Protein G purification

CloneNo.:

2H7B7

Recommended Dilutions:

WB 1:3000-1:20000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human

Positive Controls:

WB : PC-12 cells, SH-SY5Y cells

IF/ICC : SH-SY5Y cells,

Background Information

TH (Tyrosine 3-monooxygenase) converts L-tyrosine to L-3,4-dihydroxyphenylalanine (L-DOPA), the essential and rate-limiting step to formation of DA and other catecholamines. TH plays an important role in the physiology of adrenergic neurons and can be used as a marker for DA and noradrenergic neurons.

Notable Publications

Author	Pubmed ID	Journal	Application
Ting Yu	32971086	Eur J Pharmacol	WB
Amos H. P. Loh	36176417	Front Oncol	IHC
Emily-Rose Martin	36339621	Front Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of TH antibody (66334-1-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TH transfected SH-SY5Y cells.



PC-12 cells were subjected to SDS PAGE followed by western blot with 66334-1-Ig (TH antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 66334-1-Ig (TH Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using 66334-1-Ig(TH antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).