

For Research Use Only

PHF10 Monoclonal antibody

Catalog Number: 66341-1-Ig 3 Publications



Basic Information

Catalog Number: 66341-1-Ig	GenBank Accession Number: BC020954	Purification Method: Protein G purification
Size: 150ul , Concentration: 1500 ug/ml by Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 55274	CloneNo.: 4D4C6
Source: Mouse	UNIPROT ID: Q8WUB8	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Isotype: IgG1	Full Name: PHD finger protein 10	
Immunogen Catalog Number: AG19096	Calculated MW: 408 aa, 46 kDa	
	Observed MW: 56 kDa, 35 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Cited Applications: WB, IF, ColP	WB : fetal human brain tissue, HEK-293 cells, Neuro-2a cells
Species Specificity: human, mouse	IHC : human gliomas tissue,
Cited Species: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

PHF10, also named as BRG1-associated factor 45a, is a 498 amino acid protein, which locate in the nucleus and belongs to the SAYP family. PHF10 involve in transcription activity regulation by chromatin remodeling. It belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and is required for the proliferation of neural progenitors. During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth. PHF10 exists as several isoform and the calculated molecular weight of each isoform is 42 kDa, 37 kDa, 51 kDa, and 56 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Yanchen Ma	34724258	Glia	WB
Yuliang Feng	37739089	Gastroenterology	IF
Yuliang Feng	36909530	bioRxiv	ColP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



fetal human brain tissue were subjected to SDS PAGE followed by western blot with 66341-1-Ig (PHF10 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 66341-1-Ig (PHF10 Antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).