For Research Use Only

## COX2/ Cyclooxygenase 2/ PTGS2 Monoclonal antibody, PBS Only



Catalog Number:66351-1-PBS Featured Product

**Basic Information** 

Catalog Number:

GenBank Accession Number: BC013734

**Purification Method:** 

66351-1-PBS

GeneID (NCBI):

Protein A purification

Size:

100ug, Concentration: 1mg/ml by

CloneNo.: 3G2B9

Nanodrop:

**UNIPROT ID:** 

P35354

Mouse

Full Name: prostaglandin-endoperoxide synthase

Isotype: IgG2a

2 (prostaglandin G/H synthase and

Immunogen Catalog Number: AG24721

cyclooxygenase)

Calculated MW: 604 aa, 68 kDa

Observed MW:

66-70 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF-P, ELISA

Species Specificity:

human, mouse

## **Background Information**

COX2 (Prostaglandin G/H synthase 2, PTGS2) mediates the formation of prostaglandins from arachidonate. Its subunit structure is homodimer. The fully N-glycosylated PTGS2 is 72-74 kDa and the aglycosylated is 66 kDa (PMID:19656660). It also expresses a band of 39 kDa after unspecific cleavage (PMID:17509125). The 50 kDa band of fragmented PTGS2 has also previously been detected in AD brains (PMID:14724276).

Storage


Storage:

Store at -80°C.


Storage Buffer:

PBS Only


## Selected Validation Data




Non-treated and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 66351-1-Ig (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 66351-1-Ig (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66351-1-Ig (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse lung tissue using COX2/Cyclooxygenase 2/ PTGS2 antibody (66351-1-lg, Clone: 3G2B9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.