For Research Use Only

## NONO Monoclonal antibody

Catalog Number:66361-1-lg 1 Publications


Antibodies | ELISA kits | Proteins WWW.ptglab.com

Basic Information	Catalog Number: 66361-1-lg	GenBank Accession BC010049	Number:	Purification Method: Protein G purification
	Size: 150ul, Concentration: 1000 ug/ml by Bradford method using BSA as the standard; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG17609	GeneID (NCBI): 4841 UNIPROT ID: Q15233		CloneNo.: 2A2B10 Recommended Dilutions: WB 1:500-1:2000
		Calculated MW: 54 kDa Observed MW: 54 kDa		
		Applications	Tested Applications:	
, pp a constant	WB, ELISA Cited Applications: WB Species Specificity: human, rat, mouse Cited Species: mouse		WB : C2C12 ce RO51728 cells	ll, mouse skeletal muscle tissue,
Background Information	Non-pou domain-containing octamer-binding protein(NONO), other named p54nrb, belongs to DBHS family, whose protein are predominately nuclear and involves in transcriptional and posttranscriptional gene regulatory functions as well as DNA repair. This family has roles in a wide range of biological processes, such as the regulation of circadian rhythm, carcinogenesis and progression of cancer. NONO forms heterodimer with SFPQ, and has been implicated in nuclear retention of A- to I-edited RNA as paraspeckle components, pre-mRAN 3'-end formation, cAMP cycling and transcriptional activation.			
Notable Publications	Author Pub	med ID Jour	rnal	Application
	Nian Liu 359.	53472 Nat	Commun	WB
Storage	Storage: Store at -20°C. Stable for one year afte	er shipment.		
JUIGEE	Storage Buffer: PBS with 0.02% sodium azide and 50°	% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



C2C12 cell were subjected to SDS PAGE followed by western blot with 66361-1-1g (NONO Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.