For Research Use Only

Beta Catenin Monoclonal antibody, PBS Only



Purification Method:

CloneNo.:

1B8A1

Protein A purification

Catalog Number:66379-1-PBS

Featured Product

Basic Information

Catalog Number:

66379-1-PBS

100ug, Concentration: 1 mg/ml by

Nanodrop:

Source: Mouse Isotype:

lgG1

GenBank Accession Number:

NM 001904 GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000168036 **UNIPROT ID:**

P35222 Full Name:

catenin (cadherin-associated protein), beta 1, 88kDa

Calculated MW: 781 aa, 86 kDa Observed MW: 92 kDa

Applications

Tested Applications: WB, IP, IF, FC, IHC, ELISA Species Specificity: human, mouse, rat, pig

Background Information

 $\beta\text{-}Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. } \beta\text{-}Catenin$ was originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to a-catenin and the actin cytoskeleton. Besides its essential role in the AJs, β -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/β-catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β-catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222)


Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66379-1-1g (Beta Catenin antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 66379-1-lg (Beta Catenin antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 66379-1-1g (Beta Catenin antibody) at dilution of 1:20000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.



IP result of anti-Beta Catenin (IP:66379-1-Ig, 4ug; Detection:66379-1-Ig 1:2000) with mouse brain tissue lysate 4400 ug. This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Beta Catenin antibody (66379-1-lg, Clone: 1B8A1) at dilution of 1:1500 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), Alpha Tubulin antibody (11224-1-AP, red). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.



1X10^6 MCF-7 cells were intracellularly stained with 0.5 ug Anti-Human Beta Catenin (66379-1-1g, Clone:1B8A1) (red) labeled with FlexAble CoraLite® Plus 555 Antibody Labeling Kit for Mouse IgG1 (KFA022), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.