For Research Use Only

## Beta Catenin Monoclonal antibody, PBS Only



Catalog Number:66379-1-PBS

Featured Product

**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

66379-1-PBS

NM 001904 GeneID (NCBI): Protein G purification

Size:

CloneNo.: 1B8A1

100ug, Concentration: 1mg/ml by Nanodrop:

ENSEMBL Gene ID:

Source: Mouse

ENSG00000168036

Isotype: lgG1

**UNIPROT ID:** P35222 Full Name:

catenin (cadherin-associated protein), beta 1, 88kDa

Calculated MW: 781 aa, 86 kDa Observed MW:

92 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, FC (Intra), IP, ELISA

**Species Specificity:** human, mouse, rat, pig

**Background Information** 


 $\beta\text{-}Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. } \beta\text{-}Catenin$ was originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to a-catenin and the actin cytoskeleton. Besides its essential role in the AJs,  $\beta$ -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/β-catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β-catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222)

Storage


Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66379-1-1g (Beta Catenin antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 66379-1-Ig (Beta Catenin antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 66379-1-Ig (Beta Catenin antibody) at dilution of 1:20000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.




IP result of anti-Beta Catenin (IP:66379-1-Ig, 4ug; Detection:66379-1-Ig 1:2000) with mouse brain tissue lysate 4400 ug. This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Beta Catenin antibody (66379-1-lg, Clone: 188A1) at dilution of 1:1500 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), Alpha Tubulin antibody (11224-1-AP, red). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.



1X10^6 MCF-7 cells were intracellularly stained with 0.5 ug Anti-Human Beta Catenin (66379-1-lg, Clone:1B8A1) (red) labeled with FlexAble CoraLite® Plus 555 Antibody Labeling Kit for Mouse IgG1 (KFA022), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 66379-1-lg (Beta Catenin antibody) at dilution of 1:40000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human colon cancer tissue using Beta Catenin antibody (66379-1-lg, Clone: 188A1) at dilution of 1:400 and Coralite@488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66379-1-PBS in a different storage buffer formulation.