

For Research Use Only

MICA Monoclonal antibody

Catalog Number: 66384-1-Ig **3 Publications**



Basic Information

Catalog Number: 66384-1-Ig	GenBank Accession Number: BC016929	Purification Method: Protein A purification
Size: 150ul , Concentration: 1500 µg/ml by Nanodrop;	GeneID (NCBI): 100507436	CloneNo.: 1E2C8
Source: Mouse	UNIPROT ID: Q29983	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500
Isotype: IgG1	Full Name: MHC class I polypeptide-related sequence A	
Immunogen Catalog Number: AG3298	Calculated MW: 383 aa, 43 kDa	
	Observed MW: 43-50 kDa	

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HaCaT cells, Daudi cells, Raji cells, Jurkat cells, HeLa cells, A431 cells

IHC : human tonsillitis tissue, human colon tissue

Background Information

Human MHC class I chain-related genes located within the HLA class I region of chromosome 6 encode MHC class I chain-related A and B (MICA and MICB) (PMID: 11429322). MICA and MICB are stress-inducible surface molecules that are not associated with β 2-microglobulin and do not present peptides (PMID: 9497295). They are expressed in intestinal epithelium and many epithelial tumors (PMID: 10359807). MICA and MICB are ligands for NKG2D which is an activating receptor that is expressed on most natural killer (NK) cells, CD8 $\alpha\beta$ T cells, and $\gamma\delta$ T cells (PMID: 10426993, 11491531).

Notable Publications

Author	Pubmed ID	Journal	Application
Xin Fang	35183714	Pharmacol Res	IF
Chunmei Yang	38043385	Phytomedicine	WB,IF
Deli Mao	37591356	Cancer Lett	WB,IHC,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66384-1-Ig (MICA antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66384-1-Ig (MICA Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66384-1-Ig (MICA Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).