For Research Use Only

GSDMD Monoclonal antibody

Catalog Number: 66387-1-lg 41 Publications



Basic Information

Catalog Number: GenBank Accession Number:

66387-1-lg BC008904 GeneID (NCBI): Size:

150ul , Concentration: 1000 ug/ml by 79792Nanodrop: **UNIPROT ID:**

P57764

Mouse Full Name: Isotype: gasdermin D

IgG2a Calculated MW: Immunogen Catalog Number: 484 aa, 53 kDa AG16827

Observed MW: 47-53 kDa

Applications

Tested Applications:

WB, IF/ICC, IP, ELISA **Cited Applications:** WB, IHC, IF

Species Specificity: human

Cited Species: human, mouse, rat **Purification Method:**

Protein A purification

CloneNo.: 2E6C7

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:50-1:500

Positive Controls:

WB: HEK-293 cells, A431 cells, COLO 320 cells, EC109 cells, HL-60 cells, human colon tissue, human small intestine tissue, K-562 cells, Raji cells, TF-1 cells, PC-3 cells, U2OS cells, A549 cells, HepG2 cells, Jurkat cells

IP: THP-1 cells, IF/ICC: HeLa cells,

Background Information

GSDMD is a member of the gasdermin family. Members of this family appear to play a role in regulation of epithelial proliferation. GSDMD has been suggested to act as a tumor suppressor. Recently, GSDMD is identified as a new component of inflammasomes, and it is an executor of pyroptosis and required for interleukin-1β secretion. $\dot{\text{Cleavage}}$ of GSDMD by inflammatory caspases determines pyroptotic cell death.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoying Lou	36068291	Commun Biol	WB
Cintia Cevallos	36205031	J Neurochem	WB
Diego S Ojeda	30515154	Front Immunol	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66387-1-1g (GSDMD antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 66387-1-lg(GSDMD antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Mouse IgG(H+L).



IP result of anti-GSDMD (IP:66387-1-Ig, 4ug; Detection:66387-1-Ig 1:2000) with THP-1 cells lysate 1560 ug.