## For Research Use Only

## **HVCN1** Monoclonal antibody

Catalog Number: 66449-1-lg



**Basic Information** 

Catalog Number: GenBank Accession Number:

66449-1-lg BC032672 GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 1800 ug/ml by 84329 1E4C4 Nanodrop and 1000 ug/ml by Bradford<sub>UNIPROT ID:</sub> method using BSA as the standard; Q96D96

Source: Full Name: Mouse hydrogen voltage-gated channel 1

Isotype: Calculated MW: lgG1 273 aa, 32 kDa Immunogen Catalog Number: Observed MW:

28-35 kDa, 40 kDa, 60 kDa

**Purification Method:** Protein G purification

Recommended Dilutions:

WB 1:2000-1:16000 IHC 1:50-1:500 IF-P 1:200-1:800

**Applications** 

**Tested Applications:** WB, IHC, IF-P, ELISA

Species Specificity: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Raji cells, HeLa cells IHC: human tonsillitis tissue, IF-P: human tonsillitis tissue,

## **Background Information**

HVCN1, also named as VSOP and HV1, Belongs to the hydrogen channel family. HVCN1 mediates the voltagedependent proton permeability of excitable membranes. It forms a proton-selective channel through which protons may pass in accordance with their electrochemical gradient. Proton efflux, HVCN1 is accompanied by membrane depolarization, facilitates acute production of reactive oxygen species in phagocytosis. HVCN1, the voltagesensitive proton channel, is present in human sperm and is an important regulator of the functional maturation of sperm. HVCN1 has four isoforms with MW 28-32 kDa or 40 kDa (modification). It has a dimer form with MW 60 kDa.

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

E: proteintech@ptglab.com


W: ptglab.com

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Raji cells were subjected to SDS PAGE followed by western blot with 66449-1-lg (HVCN1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using HVCN1 antibody (66449-1-Ig, Clone: 1E4C4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using HVCN1 antibody (66449-1-lg, Clone: 1E4C4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66449-1-lg (HVCN1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66449-1-Ig (HVCN1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).