#### For Research Use Only

# IRF4 Monoclonal antibody

Catalog Number:66451-1-lg 2 Publications



**Purification Method:** 

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:200-1:800

**Basic Information** 

Catalog Number: GenBank Accession Number:

66451-1-Ig BC015752 Protein G purification
Size: GeneID (NCBI): CloneNo.:
150ul . Concentration: 1600 ug/ml by 3662 1A12D5

150ul , Concentration: 1600 µg/ml by 3662
Nanodrop and 1000 µg/ml by Bradford<sub>Full Name</sub>:
method using BSA as the standard;
IRF 4

 Source:
 Calculated MW:

 Mouse
 52 kDa

 Isotype:
 Observed MW:

 IgG1
 50 kDa

Immunogen Catalog Number:

AG25203

Positive Controls:

WB: human spleen tissue, pig spleen tissue

IHC: human tonsillitis,

**Applications** 

Tested Applications: IHC, WB, ELISA

Cited Applications: CoIP, IF, WB

Species Specificity: human, pig, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

## **Background Information**

IF regulatory factor 4 (IRF4) is a member of the IRF family of transcription factors, expressed in most cell types of the immune system. IRF4 is a transcription factor essential for the development of T helper-2 (Th2) cells, IL17-producing Th17 cells, and IL9-producing Th9 cells, as well as dendritic cell (DC). It binds to and activates the IF-stimulated response element (ISRE) of the MHC class I promoter, and may have a role in ISRE-targeted signal transduction mechanisms specific to lymphoid cells.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jiwei Wang	34670117	Biochim Biophys Acta Mol Cell Res	WB,CoIP
Yong Mou	35600643	Bioeng Transl Med	WB,IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### **Selected Validation Data**



human spleen tissue were subjected to SDS PAGE followed by western blot with 66451-1-1g (IRF4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66451-1-Ig (IRF4 antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66451-1-Ig (IRF4 antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).