For Research Use Only

## JAK1 Monoclonal antibody

Catalog Number:66466-1-lg Featured Product 61 Publications







Basic Information	Catalog Number: 66466-1-lg	GenBank Accession Number: BC 132729		Purification Method: Protein A purification
	Size:	GenelD (NCBI):		Protein A purification CloneNo.: 3H7A8 Recommended Dilutions:
	150ul , Concentration: 1500 ug/ml by			
	Nanodrop;			
	Source:	P23458		WB 1:1000-1:6000
	Mouse	Full Name:		IHC 1:100-1:400
	Isotype: IgG2b	Janus kinase 1 (a protein tyrosine kinase) : Calculated MW: 1154 aa, 133 kDa		
	Immunogen Catalog Number: AG17940			
		Observed MW: 133 kDa		
Applications	Tested Applications:		Positive Controls: WB : K-562 cells, Jurkat cells, rat spleen tissue, PC-12	
	WB, IHC, ELISA			
	WB, IHC, IF, IP Daudi cells		T3 cells, RAW 264.7 cells, Ramos cells,	
	Species Specificity: human, rat, mouse		IHC : human cancer tissue	preast cancer tissue, human cervical
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w	vely, antigen		
Background Information	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0 Janus kinase 1(JAK1), is a member of second phosphotransferase-related d	vely, antigen ith citrate fa class of protein-tyros omain immediately N- I cytokines. JAK1 plays on of JAK1 in cancer cell	erminal to the a critical role i s enables indi	n initiating responses to multiple maj vidual cells to contract, potentially
	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0 Janus kinase 1(JAK1), is a member of second phosphotransferase-related d signaling for certain type I and type I cytokine receptor families. Expressio allowing them to escape their tumor	vely, antigen ith citrate fa class of protein-tyros omain immediately N- I cytokines. JAK1 plays on of JAK1 in cancer cell	terminal to the a critical role i s enables indiver parts of the b	PTK domain. JAK1 is essential for n initiating responses to multiple majoridual cells to contract, potentially
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0   Janus kinase 1(JAK1), is a member of second phosphotransferase-related d signaling for certain type I and type I cytokine receptor families. Expressio allowing them to escape their tumor   Author Public	rely, antigen ith citrate	terminal to the a critical role i s enables indiver parts of the b	PTK domain. JAK1 is essential for n initiating responses to multiple maj vidual cells to contract, potentially pody.
Background Information	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0Janus kinase 1(JAK1), is a member of second phosphotransferase-related d signaling for certain type I and type I cytokine receptor families. Expression allowing them to escape their tumorAuthorPublic Shiyi LuShiyi Lu361	vely, antigen ith citrate Fa class of protein-tyros omain immediately N-I I cytokines. JAK1 plays on of JAK1 in cancer cella and metastasize to othe med ID Journa 39880 Antiox	terminal to the a critical role i s enables indiv er parts of the b	PTK domain. JAK1 is essential for in initiating responses to multiple maj vidual cells to contract, potentially body. Application

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66466-1-1g (JAK1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66466-1-1g (JAK1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 66466-1-lg (JAK1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.