

For Research Use Only

Caspase 3/p17/p19 Monoclonal antibody, PBS Only



Catalog Number: 66470-2-PBS

Featured Product

Basic Information

Catalog Number:

66470-2-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG25029

GenBank Accession Number:

BC016926

GeneID (NCBI):

836

UNIPROT ID:

P42574

Full Name:

caspase 3, apoptosis-related cysteine peptidase

Calculated MW:

277 aa, 32 kDa

Observed MW:

32-35 kDa, 19 kDa, 17 kDa

Purification Method:

Protein G purification

CloneNo.:

2G4B2

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

Human, mouse, rat

Background Information

Caspases, a family of endoproteases, are critical players in cell regulatory networks controlling inflammation and cell death. Initiator caspases (caspase-2, -8, -9, -10, -11, and -12) cleave and activate downstream effector caspases (caspase-3, -6, and -7), which in turn execute apoptosis by cleaving targeted cellular proteins. Caspase 3 (also named CPP32, SCA-1, and Apopain) proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at the beginning of apoptosis. Caspase 3 plays a key role in the activation of sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase 3 can also form heterocomplex with other proteins and performs the molecular mass of 50-70 kDa. This antibody can recognize p17, p19 and p32 of Caspase 3.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS only

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of Caspase 3 antibody (66470-2-Ig; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Caspase 3 transfected HEK-293 cells. This data was developed using the same antibody clone with 66470-2-PBS in a different storage buffer formulation.



Untreated and Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 66470-2-Ig (CASP3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66470-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66470-2-Ig (CASP3 antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66470-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66470-2-Ig (CASP3 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66470-2-PBS in a different storage buffer formulation.