

For Research Use Only

Periostin Monoclonal antibody

Catalog Number: 66491-1-Ig

Featured Product

14 Publications



Basic Information

Catalog Number: 66491-1-Ig	GenBank Accession Number: BC106710	Purification Method: Protein A purification
Size: 150ul, Concentration: 1900 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 10631	CloneNo.: 1A11A3
Source: Mouse	Full Name: periostin, osteoblast specific factor	Recommended Dilutions: WB 1:2500-1:10000 IHC 1:4000-1:16000 IF 1:4000-1:16000
Isotype: IgG1	Calculated MW: 93 kDa	
Immunogen Catalog Number: AG14487	Observed MW: 85-90 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

ELISA, FC, IF, IHC, WB

Species Specificity:

Human, Mouse, Rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : BxPC-3 cells, Neuro-2a cells, human placenta tissue, SGC-7901 cells, ROS1728 cells

IHC : human breast cancer tissue, human colon cancer tissue

IF : human breast cancer tissue,

Background Information

Periostin (POSTN, PN), originally named as osteoblast-specific factor 2 (OSF-2), is a 90-kDa secreted protein which is now classified as a matricellular protein. It is present in a wide variety of normal adult tissues and fetal tissues, and has a role in bone, tooth and heart development and function. Studies show that periostin is overexpressed in a broad range of human cancer types, including lung, ovary, breast and colon cancers. Recent evidence reveals that periostin is expressed by fibroblasts in the normal tissue and in the stroma of the primary tumour, and it is required to allow cancer stem cell maintenance. The isoforms of periostin are between 83 and 93 kDa in mass and differ in their C-terminal sequences, characterized by individual presence or absence of cassette exons 17-21 (UniProtKB/Swiss-Prot, PMID: 21997759).

Notable Publications

Author	Pubmed ID	Journal	Application
Lindsay B Alcaraz	36346290	Int J Cancer	IF
Jinna Wu	35619555	Mol Ther	WB, IHC
Jinsheng Li	35907353	Colloids Surf B Biointerfaces	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




BxPC-3, SGC-7901 and ROS1728 cells were subjected to SDS PAGE followed by western blot with 66491-1-Ig (Periostin antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66491-1-Ig (Periostin antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66491-1-Ig (Periostin antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using Periostin antibody (66491-1-Ig, Clone: 1A11A3) at dilution of 1:8000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using Periostin antibody (66491-1-Ig, Clone: 1A11A3) at dilution of 1:8000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).