For Research Use Only

OLIG2 (for WB only) Monoclonal antibody



Catalog Number:66513-1-lg

2 Publications

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: 66513-1-lg BC047511 Protein A purification

Size: GeneID (NCBI): CloneNo.: 150ul , Concentration: 1500 µg/ml by 10215 1E8C5

Nanodrop; Full Name: Recommended Dilutions:

Source: oligodendrocyte lineage transcription WB 1:1000-1:6000

Mouse factor 2
Isotype: Calculated MW:
IgG2a 32 kDa
Immunogen Catalog Number: Observed MW:
AG18838 35-39 kDa

Applications

Tested Applications: Positive Controls:

WB, ELISA WB: human brain tissue, human cerebellum tissue Cited Applications:

Species Specificity: Human

Cited Species: human

Background Information

OLIG2, also named as BHLHB1, BHLHE19, PRKCBP2 and RACK17, is required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Cooperates with OLIG1, OLIG2 establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced V3 interneuron development. OLIG2 is widely expressed in subsets of glia cells and progenitors, and it is strongly induced at different sites by both acute and chronic injury, albeit with different mechanisms. OLIG2 acts as a repressor of neurogenesis in cells reacting to brain injury. It may represent an effective approach towards evoking neuronal repair from parenchymal precursors.(PMID:19390819)

Notable Publications

Author	Pubmed ID	Journal	Application
Hongtao Zhu	35814469	Front Oncol	WB
Xianli Du	36660824	Nucleic Acids Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 66513-1-1g (OLIG2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.