

For Research Use Only

IGF2BP3 Monoclonal antibody

Catalog Number: 66526-1-Ig **1 Publications**



Basic Information

Catalog Number: 66526-1-Ig	GenBank Accession Number: BC065269	Purification Method: Protein G purification
Size: 150ul , Concentration: 747 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 10643	CloneNo.: 2C10B11
Source: Mouse	UNIPROT ID: O00425	Recommended Dilutions: WB 1:1000-1:6000
Isotype: IgG1	Full Name: insulin-like growth factor 2 mRNA binding protein 3	
Immunogen Catalog Number: AG6349	Calculated MW: 64 kDa	
	Observed MW: 65 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HSC-T6 cells, NIH/3T3 cells, HEK-293 cells, Jurkat cells, HepG2 cells, K-562 cells
Cited Applications: WB	
Species Specificity: Human, mouse, rat	
Cited Species: human	

Background Information

IGF2BP3, also named as IMP3, KOC1 and VICKZ3, belongs to the RRM IMP/VICKZ family. It is one of the RNA binding proteins involved in mRNA localization and translational control. IGF2BP3 is expressed during embryogenesis, as well as in some malignant tumors. It can be used as an independent prognostic factor for osteosarcoma.

Notable Publications

Author	Pubmed ID	Journal	Application
Luis Daniel Cruz-Zaragoza	34672953	Cell	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various cells were subjected to SDS PAGE followed by western blot with 66526-1-Ig (IGF2BP3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66526-1-Ig (IGF2BP3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.