For Research Use Only

Cathepsin D Monoclonal antibody

Catalog Number:66534-1-lg Featured Product

7 Publications

32 kDa, 48 kDa, 52 kDa



Basic Information

Catalog Number: GenBank Accession Number:

66534-1-lg BC016320 GeneID (NCBI):

150ul, Concentration: 1000 µg/ml by 1509

Full Name: Source: cathepsin D Mouse Calculated MW: Isotype: 412 aa, 45 kDa lgG2b Observed MW:

AG15254

Purification Method: Protein A purification

CloneNo.:

2F6F7

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:500-1:2000 IF 1:200-1:800

Applications

Tested Applications:

Immunogen Catalog Number:

FC, IF, IHC, WB, ELISA Cited Applications:

IF, IHC, WB

Species Specificity:

Human **Cited Species:** human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Saos-2 cells, MCF-7 cells, T-47D cells, HepG2 cells

IHC: human liver cancer tissue, human breast cancer

tissue

IF: HepG2 cells,

Background Information

CTSD (Cathepsin D) is also named as CPSD, belongs to the peptidase A1 family. It is ubiquitously expressed and is involved in proteolytic degradation, cell invasion, and apoptosis. Human CTSD is synthesized as a 52-kDa precursor that is converted into an active 48-kDa single-chain intermediate in the endosomes, and then into a fully active mature form, composed of a 34-kDa heavy chain and a 14-kDa light chain, in the lysosomes. It is a lysosomal acid protease found in neutrophils and monocytes and involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease. (PMID: 27114232, PMID: 30717773, PMID: 30051532)

Notable Publications

Author	Pubmed ID	Journal	Application
Zhiyuan Wu	34575099	Life (Basel)	WB,IF
Hualin Fu	36448495	FEBS Lett	IF
Peipei Ding	35649359	Cell Rep	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human Cathepsin D (66534-1-1g, Clone:2F6F7) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-1g, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).




Various lysates were subjected to SDS PAGE followed by western blot with 66534-1-1g (Cathepsin D antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66534-1-1g (Cathepsin D antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66534-1-Ig (Cathepsin D antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Cathepsin D antibody (66534-1-Ig, Clone: 2F6F7) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).