For Research Use Only

TBR1 Monoclonal antibody

Catalog Number:66564-1-lg 5 Publications



Basic Information

Catalog Number: GenBank Accession Number:

66564-1-lg BC104844 GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 2000 µg/ml by 10716 5C3A5 Nanodrop and 1069 µg/ml by Bradford_{Full Name}:

T-box, brain, 1

Calculated MW: Mouse 682 aa, 74 kDa

Isotype: Observed MW: lgG1 74 kDa

Immunogen Catalog Number:

method using BSA as the standard;

AG15262

Applications

Tested Applications:

IF, IHC, WB, ELISA **Cited Applications:**

IF, IHC

Species Specificity: Human, rat, mouse, pig

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Protein G purification

Recommended Dilutions: WB 1:2000-1:12000 IHC 1:50-1:500 IF 1:50-1:500

Positive Controls:

WB: pig brain tissue, pig cerebellum tissue, rat cerebellum tissue, rat brain tissue, mouse brain tissue,

rabbit brain tissue IHC: mouse brain tissue.

IF: mouse brain tissue.

Background Information

TBR1, also named T-box brain protein 1, is a 682 amino acid protein, which contains one T-box DNA-binding domain and localizes in the nucleus. TBR1 is expressed in the brain and as a transcriptional regulator is involved in developmental processes. TBR1 is required for normal brain development.

Notable Publications

Author	Pubmed ID	Journal	Application
Thao P Phan	33226141	EMBO J	IHC
Egidio Spinelli	34185323	Ann Neurol	IF
Huai-Bin Hu	33510165	Nat Commun	IF

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66564-1-Ig (TBR1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66564-1g (TBR1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66564-1-Ig (TBR1 antibody) at dilution of 1:200 (under 4x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 66564-1-lg (TBR1 antibody) at dilution of 1:100 and Alexa Fluor 594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 66564-1-lg (TBR1 antibody) at dilution of 1:100 and Alexa Fluor 594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



pig cerebellum tissue were subjected to SDS PAGE followed by western blot with 66564-1-1g (TBR1 antibody) at dilution of 1:2000 incubated at 4 degree celsius over night.