

For Research Use Only

# NXN Monoclonal antibody

Catalog Number: 66603-1-Ig



## Basic Information

<b>Catalog Number:</b> 66603-1-Ig	<b>GenBank Accession Number:</b> BC009327	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 150ul , Concentration: 2100 ug/ml by Nanodrop and 1000 ug/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 64359	<b>CloneNo.:</b> 2C6A4
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q6DKJ4	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IHC 1:50-1:500 IF/ICC 1:400-1:1600
<b>Isotype:</b> IgG1	<b>Full Name:</b> nucleoredoxin	
<b>Immunogen Catalog Number:</b> AG9248	<b>Calculated MW:</b> 435 aa, 48 kDa	
	<b>Observed MW:</b> 48 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IF/ICC, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> Human, Mouse, Rat	<b>WB :</b> HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells
<b>Note-IHC:</b> suggested antigen retrieval with <b>TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	<b>IHC :</b> human kidney tissue,
	<b>IF/ICC :</b> HEK-293 cells,

## Background Information

Nucleoredoxin (NRX/NXN), a relatively uncharacterized member of the TRX family protein, has recently been reported to regulate the Wnt/ $\beta$ -catenin pathway, which itself regulates cell fate and early development, in a redox-dependent manner (PMID:17567240). Northern blot analysis detected several nucleoredoxin splice variants in adult mouse tissues, with highest expression in testis and skin and lowest expression in liver (PMID:9119370). It has 3 isoforms produced by alternative splicing. This antibody is specific to NXN.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using NXN antibody (66603-1-Ig, Clone: 2C6A4 ) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 66603-1-Ig (NXN antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 66603-1-Ig (NXN antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66603-1-Ig (NXN antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.