#### For Research Use Only

# Adiponectin receptor 1 Monoclonal antibody



Catalog Number: 66619-1-Ig

**Featured Product** 

6 Publications

**Basic Information** 

Catalog Number: 66619-1-lg

GenBank Accession Number: BC010743

Protein G purification

WB 1:1000-1:6000

**Purification Method:** 

Size:

GeneID (NCBI):

CloneNo.:

150ul, Concentration: 1000 µg/ml by 51094 Bradford method using BSA as the

1F12B3

adiponectin receptor 1

Recommended Dilutions:

standard:

Calculated MW:

Mouse Isotype:

375 aa, 43 kDa

Observed MW:

lgG1

AG18187

Immunogen Catalog Number:

43 kDa

**Positive Controls:** 

**Applications** 

**Tested Applications:** 

WB,ELISA

WB: HeLa cells, HEK-293 cells, LO2 cells, HepG2 cells

**Cited Applications:** 

Species Specificity:

Human

**Cited Species:** 

human, mouse, pig

### **Background Information**

Adiponectin is a hormone secreted by adipocytes that acts as an antidiabetic and anti-atherogenic adipokine. Two receptors for adiponectin (ADIPOR1 and ADIPOR2) have been identified. They mediate increased AMPK, MAPK and PPARA ligand activity in response to adiponectin. AdipoR1 is abundantly expressed in skeletal muscle, whereas AdipoR2 is predominantly expressed in the liver. The human ADIPOR1 gene maps to chromosome 1q32.1, and encodes a 375-amino acid protein with a molecular weight of 43 kDa. This antibody was raised against 1-72aa of human AdipoR1.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaoling Chen	34543141	Anim Biotechnol	WB
Xin Li	34019043	Food Funct	WB
Lu Xiang	33703997	Anim Biotechnol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data





WB result of Adiponectin receptor 1 antibody (66619-1-lg; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Adiponectin receptor 1 transfected HEK-293 cells.

HeLa cells, LO2 cells, and HepG2 cells were subjected to SDS PAGE followed by western blot with 66619-1-lg (Adiponectin receptor 1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.