For Research Use Only

FAF2 Monoclonal antibody

Catalog Number:66629-1-lg Featured Product



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

66629-1-lg BC014001 Protein G purification CloneNo.:

GeneID (NCBI): Size: 150ul, Concentration: 1700 ug/ml by 23197 3H6D2

Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} Recommended Dilutions: method using BSA as the standard; Q96CS3 WB 1:2000-1:10000 Source: IHC 1:250-1:1000 Full Name:

Mouse Fas associated factor family member

Isotype:

lgG1 Calculated MW: Immunogen Catalog Number: 445 aa. 53 kDa AG21536 Observed MW:

51 kDa

Applications

Tested Applications: Positive Controls:

WB, IHC, ELISA WB: HeLa cells, HEK-293 cells, HepG2 cells

Species Specificity: IHC: human breast cancer tissue, mouse brain tissue Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Protein ETEA (FAF2, or UBXD8) is a homolog to Fas-associated factor 1 (FAF1), which is involved in Fas-mediated apoptosis. ETEA protein directly interacts with and negatively regulates neurofibromin. ETEA contains both UBA and UBX domains and overexpression of ETEA downregulates neurofibromin in human cells. It may play a role in the translocation of terminally misfolded proteins from the endoplasmic reticulum lumen to the cytoplasm and their degradation by the proteasome. ETEA is highly expressed in peripheral blood of patients with atopic dermatitis (AD) $compared \ to \ normal \ individuals, and \ may \ regulate \ the \ resistance \ to \ apoptosis \ that \ is \ observed \ in \ T \ cells \ and$ eosinophils of AD patients.

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 66629-1-lg (FAF2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66629-1-1g (FAF2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of FAF2 antibody (66629-1-lg; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FAF2 transfected HeLa cells.