

For Research Use Only

IRAK1 Monoclonal antibody

Catalog Number: 66653-1-Ig

Featured Product

1 Publications



Basic Information

Catalog Number: 66653-1-Ig	GenBank Accession Number: BC014963	Purification Method: Protein A purification
Size: 150UL, Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 3654	CloneNo.: 1H10A7
Source: Mouse	Full Name: interleukin-1 receptor-associated kinase 1	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:250-1:1000
Isotype: IgG2a	Calculated MW: 77 kDa	
Immunogen Catalog Number: AG0728	Observed MW: 80 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IF

Species Specificity:

Human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells

IHC : human lung cancer tissue,

Background Information

Interleukin-1 receptor-associated kinases (IRAKs) are a unique family of death domain containing protein kinases that play a key role in initiating innate immune response against foreign pathogens. They are involved in Toll-like receptor (TLR) and interleukin-1 receptor (IL-1R) signaling pathways. IRAK1 is the first member of this kinase family. Upon ligand binding to TLR/IL-1R, IRAK1 is recruited by MYD88 to the receptor-signaling complex, the association leads to IRAK1 phosphorylation by IRAK4 and subsequent autophosphorylation and kinase activation. Hyper-phosphorylated IRAK1 then disengages from the receptor complex, and forms a cytosolic IRAK1-TRAF6 complex. TRAF6 then interacts with TAK and TAB, resulting in eventual activation of the NF- κ B and MAPK pathways. Phosphorylated IRAK1 also undergoes ubiquitin-mediated degradation or sumoylation, which results in nuclear translocation and transcriptional activation of inflammatory target genes. (PMID: 17890055; 12620219)

Notable Publications

Author	Pubmed ID	Journal	Application
Gang Xu	33664485	Cell Mol Immunol	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of IRAK1 antibody (66653-1-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IRAK1 transfected HEK-293 cells.



HeLa cells were subjected to SDS PAGE followed by western blot with 66653-1-Ig (IRAK1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66653-1-Ig (IRAK1 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66653-1-Ig (IRAK1 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).