For Research Use Only

PSMA/GCPII Monoclonal antibody

Catalog Number:66678-1-lg 1 Publications



Purification Method:

Recommended Dilutions:

WB 1:4000-1:10000

IHC 1:1000-1:4000 IF 1:200-1:800

Basic Information

Catalog Number: GenBank Accession Number:

66678-1-lg Protein A purification BC025672 GeneID (NCBI): CloneNo.:

150ul, Concentration: 1000 µg/ml by 2346 3G4F12 Bradford method using BSA as the

standard: folate hydrolase (prostate-specific

membrane antigen) 1 Mouse Calculated MW: 719 aa, 81 kDa Isotype: lgG1

Observed MW: 100-120 kDa Immunogen Catalog Number:

AG16594

Tested Applications: Applications IF THE WRIFTISA

Cited Applications:

Species Specificity: Human, Rat **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, rat kidney tissue IHC: human prostate cancer tissue,

IF: human prostate cancer tissue, PC-3 cells

Background Information

PSMA(Prostate-specific membrane antigen) is also named as FOLH1, FOLH, NAALAD1, PSM and belongs to the $peptidase\ M28\ family.\ PSMA\ is\ a\ 100-120\ kDa\ integral\ transmembrane\ glycoprotein,\ considered\ to\ be\ a\ highly$ specific marker of the prostate gland, and has successfully been used as a marker of circulating prostatic epithelial cells(PMID:10074909; 15680901). It is involved in conversion of the major neurotransmitter (NAAG) to NAA and free glutamate. It has 8 isoforms produced by alternative splicing.

Notable Publications

Author **Pubmed ID** Journal Application 31988304 WB Liyuan Yin Sci Ren

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66678-1-Ig (PSMA/GCPII antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



LNCaP cells were subjected to SDS PAGE followed by western blot with 66678-1-Ig (FOLH1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66678-1-Ig (PSMA/GCPII antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human prostate cancer tissue using PSMA/GCPII antibody (66678-1-Ig, Clone: 3G4E12) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).