

UCP2 Monoclonal antibody, PBS Only

Catalog Number: 66700-1-PBS

Basic Information

Catalog Number:	66700-1-PBS	GenBank Accession Number:	BC011737	Purification Method:	Protein A purification
Size:	100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI):	7351	CloneNo.:	2B4B1
Source:	Mouse	UNIPROT ID:	P55851		
Isotype:	IgG3	Full Name:	uncoupling protein 2 (mitochondrial, proton carrier)		
Immunogen Catalog Number:	AG19361	Calculated MW:	33 kDa		
		Observed MW:	28 kDa		

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
Human

Background Information

UCP2 (uncoupling protein 2) is a member of the mitochondrial carrier protein super family located in the mitochondrial inner membrane. The UCPs differ greatly in tissue distribution: UCP1 is expressed only in brown adipose; UCP3 preferentially in skeletal muscle, brown adipose and heart; UCP4 and UCP5 mainly in the nervous system (18167556). In contrast, UCP2 has a broad distribution and is implicated in a wide range of pathophysiological processes, including diabetes and cancer. This antibody detect a band around 28-33 kDa corresponding to UCP2 protein in mouse liver, skeletal muscle and pancreas. Some upper and lower bands also were observed due to the dimer formation and degradation of the UCP2 protein (20103532). This antibody was raised against the full-length UCP2 protein and may cross-react with UCP3. It is notable that UCP2 is known to have a very short half-life less than 1 hour (17240372). It is strongly recommended that fresh-made lysate be used for western blot assays.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 66700-1-Ig (UCP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66700-1-PBS in a different storage buffer formulation.



human adipose tissue were subjected to SDS PAGE followed by western blot with 66700-1-Ig (UCP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66700-1-PBS in a different storage buffer formulation.



human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 66700-1-Ig (UCP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66700-1-PBS in a different storage buffer formulation.