

For Research Use Only

MTAP Monoclonal antibody

Catalog Number: 66706-1-Ig



Basic Information

Catalog Number: 66706-1-Ig	GenBank Accession Number: BC018625	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop and 437 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 4507	CloneNo.: 2B1G6
Source: Mouse	UNIPROT ID: Q13126	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500 IF/ICC 1:400-1:1600
Isotype: IgG1	Full Name: methylthioadenosine phosphorylase	
Immunogen Catalog Number: AG17954	Calculated MW: 283 aa, 31 kDa	
	Observed MW: 31 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, ELISA

Species Specificity:
Human, Mouse, Rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : NIH/3T3 cells, HCT 116 cells, HSC-T6 cells, Raji cells, HT-29 cells

IHC : mouse liver tissue, human kidney tissue, rat liver tissue, mouse kidney tissue, human colon cancer tissue, human liver cancer tissue

IF/ICC : HeLa cells,

Background Information

MTAP is a 5-deoxy-5-methylthioadenosine (MTA) phosphorylase, converting MTA to salvageable intermediates including adenine and 5-methylthioribose-1-phosphate. MTAP is abundant in normal cells, but it is frequently found to be deleted in a variety of cancers. Its deficiency is common in cancer cell lines as well as in primary leukemia, lung cancer, melanoma, bladder cancer, gliomas and breast cancer.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66706-1-Ig (MTAP antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 66706-1-Ig (MTAP antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MTAP antibody (66706-1-Ig, Clone: 2B1G6) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).