For Research Use Only

STRAP Monoclonal antibody, PBS Only



Catalog Number:66712-1-PBS

Featured Product

Basic Information

Catalog Number: 66712-1-PBS Size: 100ug , Concentration: 1 mg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG13085 GenBank Accession Number: BC000162 GeneID (NCBI): 11171 UNIPROT ID: Q9Y3F4 Full Name: serine/threonine kinase receptor associated protein Calculated MW: 38 kDa Observed MW: 38 kDa

Purification Method: Protein G purification CloneNo.: 5E11F3

Applications

Tested Applications: WB, IF, IHC, Indirect ELISA Species Specificity: Human, mouse, rat

Background Information

STRAP (Serine-threonine kinase receptor-associated protein), also known as MAWD, is a ubiquitous WD40 domain protein, common function of which is to provide a suitable scaffold for coordinating multiprotein complex assemblies and thus, regulate a variety of cellular processes like signal transduction, transcriptional regulation, programmed cell death and so on. STRAP plays a role in the cellular distribution of the SMN complex, which is essential for spliceosomal snRNP assembly in the cytoplasm. STRAP inhibits transforming growth factor-beta (TGF-beta) signaling and enhances tumorigenicity via TGF-beta-dependent and -independent mechanisms. STRAP is up-regulated in several cancers and functions as an oncogene. STRAP regulates c-Jun stability by decreasing the ubiquitylation and proteosomal degradation of c-Jun.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66712-1-1g (STRAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66712-1-PBS in a different storage buffer formulation.




WB result of STRAP antibody (66712-1-lg; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STRAP transfected HEK-293 cells. This data was developed using the same antibody clone with 66712-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 66712-1-Ig (STRAP antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66712-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 66712-1-Ig (STRAP antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66712-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using STRAP antibody (66712-1-lg, Clone: 5E11F3) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66712-1-PBS in a different storage buffer formulation.