For Research Use Only

HSPH1 Monoclonal antibody

Catalog Number: 66723-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

66723-1-lg BC037553 Protein G purification

GeneID (NCBI): Size: CloneNo.: 150ul , Concentration: 1700 ug/ml by 10808 3H5A5

Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} Recommended Dilutions: method using BSA as the standard; Q92598 WB 1:5000-1:20000 Source: IHC 1:500-1:2000 Full Name: heat shock 105kDa/110kDa protein 1 IF/ICC 1:50-1:500 Mouse

Isotype: Calculated MW: lgG1 858 aa, 97 kDa Immunogen Catalog Number: Observed MW: AG27521 110 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Species Specificity: Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells

Purification Method:

IHC: mouse testis tissue, IF/ICC: HeLa cells,

Background Information

HSP105, also known as HSP110 or HSPH1, belongs to the heat shock protein (HSP) family. Human HSP105 is a highmolecular-weight chaperone protein expressed at constitutively low levels as a cytoplasmic α-isoform and as an inducible nuclear β-isoform on exposure to various forms of stress. HSP105 is constitutively overexpressed in several solid tumors, including melanoma, breast, thyroid, and gastroenteric cancers, and exerts antiapoptotic functions. Recently HSP105 has been identified as a novel candidate biomarker of lymphoma aggressiveness. Western blot analysis using this antibody detected the phosphorylated protein around 110 kDa in various lysates.

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66723-1-1g (HSPH1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 66723-1-Ig (HSPH1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 66723-1-Ig (HSPH1 antibody) at dilution of 1:100 and Alexa Fluor 488-Conjugated Goat Anti-Mouse IgG(H+L).