

For Research Use Only

TYMS Monoclonal antibody, PBS Only

Catalog Number: 66725-1-PBS

Featured Product



Basic Information

Catalog Number:

66725-1-PBS

Size:

100ug, Concentration: 1mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG7027

GenBank Accession Number:

BC002567

GeneID (NCBI):

7298

UNIPROT ID:

P04818

Full Name:

thymidylate synthetase

Calculated MW:

36 kDa

Observed MW:

36 kDa

Purification Method:

Protein A purification

CloneNo.:

1F5B7

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Background Information

Thymidylate synthase gene (TYMS) encodes thymidylate synthase (TS) which is an important factor in the growth of tumor cells. TS can catalyze the transformation of intracellular uridine monophosphate (UMP) into dTMP which is required for DNA replication and repairing. TS is a key enzyme in the process of cell proliferation, also is the important target enzymes of 5-FU and other chemotherapy drugs. The expression of TYMS is negatively correlated with the efficacy of chemotherapy and the prognosis of patients who suffer from rectal cancer, breast cancer, colorectal cancer, gastric cancer, head and neck cancer, esophageal cancer, pancreatic cancer and so on.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66725-1-Ig (TYMS antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.




WB result of TYMS antibody (66725-1-Ig; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TYMS transfected HEK-293 cells. This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66725-1-Ig (TYMS antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 66725-1-Ig (TYMS antibody) at dilution of 1:400 (under 10x lens). Proteolytic pre-treatment mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 66725-1-Ig (TYMS antibody) at dilution of 1:400 (under 40x lens). Proteolytic pre-treatment mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66725-1-Ig (TYMS antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using TYMS antibody (66725-1-Ig, Clone: 1F5B7) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66725-1-PBS in a different storage buffer formulation.



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug TYMS Monoclonal antibody (66725-1-Ig, Clone:1F5B7) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.25 ug Mouse IgG1 isotype control Mouse McAb (66360-1-Ig, Clone: 1F8D3) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66725-1-PBS in a different storage

